

Available online at www.sciencedirect.com

J. Math. Anal. Appl. 326 (2007) 1452–1457

Journal of
 MATHEMATICAL
 ANALYSIS AND
 APPLICATIONS

www.elsevier.com/locate/jmaa

Note

On new generalizations of the Hilbert integral inequality

Lü Zhongxue^a, Gao Mingzhe^b, Lokenath Debnath^{c,*}^a Department of Basic Science of Technology College, Xuzhou Normal University, Jiangsu 221011, China^b Department of Mathematics, Normal College, Jishou University, Hunan Jishou 416000, China^c Department of Mathematics, University of Texas – Pan American, 1201 W. University Drive, Edinburg, TX 78539, USA

Received 28 February 2006

Available online 18 April 2006

Submitted by William F. Ames

Abstract

Some new generalizations of the Hilbert integral inequality by introducing real functions $\phi(x)$ and $\psi(x)$. The results of this paper reduce to those of the corresponding inequalities proved by Gao [Mingzhe Gao, On Hilbert's integral inequality, Math. Appl. 11 (3) (1998) 32–35]. Some applications are considered.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Inner product; Hilbert integral inequality; Hardy–Littlewood integral inequality

1. Introduction

Let $f(x), g(x) \in L^2(0, \infty)$. Then Hilbert's integral inequality may be written as

$$\left(\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy \right)^2 \leq \pi^2 \int_0^\infty f^2(x) dx \int_0^\infty g^2(y) dy, \quad (1)$$

where π is the best value started by Hardy et al. [1, Chapter 9].

* Corresponding author.

E-mail addresses: lvzxl@163.net (Z. Lü), mingzhgao@163.com (M. Gao), debnathl@panam.edu (L. Debnath).

Recently, Gao [2] gave an improvement of (1) by means of Gram's matrix as follows:

$$\left(\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy \right)^2 \leq \pi^2 \int_0^\infty f^2(x) dx \int_0^\infty g^2(y) dy - F(s, t), \quad (2)$$

where the function F is defined by

$$F(s, t) = \|\alpha\|^2 s^2 - 2(\alpha, \beta)st + \|\beta\|^2 t^2, \quad (3)$$

where α , β and γ are three arbitrary vectors of inner product space E , (α, β) indicates the inner product of the vectors α and β , and γ is a unit-vector, the α and β are not simultaneously orthogonal to the vector γ , and $s = (\beta, \gamma)$ and $t = (\alpha, \gamma)$ are the functionals of the vector γ of E .

Based on the method of Gao [2], we further make some new generalizations of Hilbert integral inequality by introducing real functions $\phi(x)$ and $\psi(x)$. Some Hilbert type integral inequalities are obtained. Some applications are also considered.

2. Main results

First we give some lemmas.

Lemma 2.1. (See Gao [2].) *Let $F(s, t)$ be the function defined by (3). If the vectors α and β are linearly independent, then $F(s, t) > 0$ for any $s, t \in \mathbb{R}$, provided that s and t are not zero simultaneously.*

Lemma 2.2. (See Gao [2].) *Let α and β be two given vectors of E , and γ be an arbitrary vector of E . If the vectors α and β are linearly independent, and $\|\gamma\| = 1$, then*

$$(\alpha, \beta)^2 \leq \|\alpha\|^2 \|\beta\|^2 - F(s, t), \quad (4)$$

where $F(s, t)$ is the function defined by (3) and $F(s, t) > 0$. And the equality contained in (4) holds if and only if the vector γ is a linear combination of the vectors α and β .

Theorem 2.1. *Let $f, g \in L^2(0, \infty)$ and $\phi(x)$ and $\psi(x)$ be differentiable functions in $(0, +\infty)$, and $\phi(0), \psi(0) \geq 0$, $\phi(\infty) = \infty$, $\psi(\infty) = \infty$, $\phi(0) = \psi(0)$, $\phi'(x), \psi'(x) > 0$, $\phi'(x)$ and $\psi'(x)$ have positive infimum, respectively, then*

$$\begin{aligned} & \left(\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{\phi(x) + \psi(y)} dx dy \right)^2 \\ & \leq \frac{1}{\inf\{\phi'(x)\} \inf\{\psi'(y)\}} \int_0^\infty \left(\pi - \arctan \sqrt{\frac{\psi(0)}{\phi(x)}} \right) f^2(x) dx \\ & \quad \times \int_0^\infty \left(\pi - \arctan \sqrt{\frac{\phi(0)}{\psi(y)}} \right) g^2(y) dy - F(s, t), \end{aligned} \quad (5)$$

where $F(s, t)$ is the function defined by (3) and $F(s, t) > 0$.

Proof. We define two functions by

$$\alpha = \frac{f(x)}{(\phi(x) + \psi(y))^{1/2}} \left(\frac{\phi(x)}{\psi(y)} \right)^{1/4}, \quad \beta = \frac{g(y)}{(\phi(x) + \psi(y))^{1/2}} \left(\frac{\psi(y)}{\phi(x)} \right)^{1/4}. \quad (6)$$

We may apply inequality (4) to estimate the left-hand side of (5) as follows:

$$\begin{aligned} & \left(\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{\phi(x) + \psi(y)} dx dy \right)^2 \\ &= \left(\int_0^\infty \int_0^\infty \alpha \beta dx dy \right)^2 \\ &\leq \int_0^\infty \int_0^\infty \alpha^2 dx dy \int_0^\infty \int_0^\infty \beta^2 dx dy - F(s, t) \\ &= \int_0^\infty \left(\int_0^\infty \frac{1}{\phi(x) + \psi(y)} \left(\frac{\phi(x)}{\psi(y)} \right)^{1/2} dy \right) f^2(x) dx \\ &\quad \times \int_0^\infty \left(\int_0^\infty \frac{1}{\phi(x) + \psi(y)} \left(\frac{\psi(y)}{\phi(x)} \right)^{1/2} dx \right) g^2(y) dy - F(s, t) \\ &= \int_0^\infty \omega(\phi, \psi, x) f^2(x) dx \int_0^\infty \omega(\psi, \phi, y) g^2(y) dy - F(s, t), \end{aligned} \quad (7)$$

where the weight function $\omega(\phi, \psi, x)$ is defined by

$$\omega(\phi, \psi, x) = \int_0^\infty \frac{1}{\phi(x) + \psi(y)} \left(\frac{\phi(x)}{\psi(y)} \right)^{1/2} dy.$$

Putting $u = \frac{\psi(y)}{\phi(x)}$, we have

$$\begin{aligned} \omega(\phi, \psi, x) &\leq \frac{1}{\inf\{\psi'(y)\}} \int_{\frac{\psi(0)}{\phi(x)}}^\infty \frac{1}{1+u} \left(\frac{1}{u} \right)^{1/2} du \\ &= \frac{1}{\inf\{\psi'(y)\}} \left(\pi - 2 \arctan \sqrt{\frac{\psi(0)}{\phi(x)}} \right). \end{aligned} \quad (8)$$

Similarly we prove

$$\begin{aligned} \omega(\psi, \phi, y) &\leq \frac{1}{\inf\{\phi'(x)\}} \int_{\frac{\phi(0)}{\psi(y)}}^\infty \frac{1}{1+u} \left(\frac{1}{u} \right)^{1/2} du \\ &= \frac{1}{\inf\{\phi'(x)\}} \left(\pi - 2 \arctan \sqrt{\frac{\phi(0)}{\psi(y)}} \right). \end{aligned} \quad (9)$$

It follows from (7)–(9) that inequality (5) remains valid. Obviously, α and β defined by (6) are independent, by Lemma 2.1 we have $F(s, t) > 0$. And the theorem is proved. \square

Corollary 2.1. Let $f \in L^2(0, \infty)$ and $\phi(x)$ be a differentiable function in $(0, \infty)$. If $\phi(0) \leq 0$, $\phi(\infty) = \infty$, $\phi'(x) > 0$ and $\phi'(x)$ has a positive infimum in $(0, \infty)$, then

$$\left(\int_0^\infty \int_0^\infty \frac{f(x)f(y)}{\phi(x) + \phi(y)} dx dy \right)^2 \leq \frac{1}{\inf\{\phi'(x)\}^2} \left(\int_0^\infty \left(\pi - 2 \arctan \sqrt{\frac{\phi(0)}{\phi(x)}} \right) f^2(x) dx \right)^2 - F(s, t), \quad (10)$$

where $F(s, t)$ is the function defined by (3) and $F(s, t) > 0$.

Remark 2.1. When $\phi(x) = \psi(x) = x$, inequalities (5) and (10) reduce to (5) and (7) of the paper by Gao [2], respectively, hence inequalities (5) and (10) in this paper are generalizations of the corresponding results of Gao [2].

3. Some applications

We take $\phi(x)$ and $\psi(x)$ as

$$\phi(x) = e^x, \quad \psi(y) = e^y,$$

then, by Theorem 2.1, we get the following result:

Theorem 3.1. Let $f, g \in L^2(0, \infty)$. Then

$$\begin{aligned} \left(\int_0^\infty \int_0^\infty \frac{f(x)g(y)}{e^x + e^y} dx dy \right)^2 &\leq \int_0^\infty \left(\pi - 2 \arctan \sqrt{\frac{1}{e^x}} \right) f^2(x) dx \\ &\quad \times \int_0^\infty \left(\pi - 2 \arctan \sqrt{\frac{1}{e^y}} \right) g^2(y) dy - F(s, t), \end{aligned} \quad (11)$$

where $F(s, t)$ is defined by (3) and $F(s, t) > 0$.

Proof. We define two functions by

$$\alpha = \frac{f(x)}{(e^x + e^y)^{1/2}} e^{\frac{x-y}{4}}, \quad \beta = \frac{g(y)}{(e^x + e^y)^{1/2}} e^{\frac{y-x}{4}}. \quad (12)$$

A proof similar to that of Theorem 2.1 can be used to obtain inequality (11). We omit the proof. \square

Theorem 3.2. Let $g \in L^2(0, 1)$ with $g(t) \neq 0$ for all $t \in (0, 1)$, $\phi(x)$ be a differentiable function in $(0, \infty)$, and $\phi(0) \geq 0$, $\phi(\infty) = \infty$, $\phi'(x) > 0$ and $\phi'(x)$ has a positive infimum, and define a function F by

$$F(x) = \int_0^1 u^{\phi(x)} g(u) du \quad (x \geq 0),$$

then

$$\begin{aligned} & \left(\int_0^\infty f^2(x) dx \right)^2 \\ & \leq \left\{ \left(\frac{1}{(\inf\{\phi'(x)\})^2} \int_0^\infty \left(\pi - 2 \arctan \sqrt{\frac{\phi(0)}{\phi(x)}} \right) f^2(x) dx \right)^2 - F_2(s, t) \right\}^{1/2} \\ & \quad \times \int_0^1 u g^2(u) du - F_1(s, t), \end{aligned} \quad (13)$$

where $F_1(s, t)$, $F_2(s, t)$ are defined by (3) and $F_1(s, t) > 0$, $F_2(s, t) > 0$.

Proof. We may write $f^2(x)$ in the form

$$f^2(x) = \int_0^1 f(x) u^{\phi(x)} g(u) du.$$

We next apply inequalities (4) and (5) successively to estimate the left-hand side of (13) as follows:

$$\begin{aligned} & \left(\int_0^\infty f^2(x) dx \right)^2 \\ & = \left\{ \int_0^\infty \left(\int_0^1 f(x) u^{\phi(x)} g(u) du \right) dx \right\}^2 \\ & = \left\{ \int_0^1 \left(\int_0^\infty f(x) u^{\phi(x)-1/2} dx \right) (u^{1/2} g(u)) du \right\}^2 \\ & \leq \int_0^1 \left(\int_0^\infty f(x) u^{\phi(x)-1/2} dx \right)^2 du \int_0^1 u g^2(u) du - F_1(s, t) \\ & = \int_0^1 \left(\int_0^\infty f(x) u^{\phi(x)-1/2} dx \right) \left(\int_0^\infty f(y) u^{\phi(y)-1/2} dy \right) du \int_0^1 u g^2(u) du - F_1(s, t) \end{aligned}$$

$$\begin{aligned}
&= \int_0^1 \left(\int_0^\infty \int_0^\infty f(x) f(y) u^{\phi(x)+\phi(y)-1} dx dy \right) du \int_0^1 u g^2(u) du - F_1(s, t) \\
&= \int_0^\infty \int_0^\infty \frac{f(x) f(y)}{\phi(x) + \psi(y)} dx dy \int_0^1 u g^2(u) du - F_1(x, y) \\
&\leq \left\{ \left(\frac{1}{(\inf\{\phi'(x)\})^2} \int_0^\infty \left(\pi - 2 \arctan \sqrt{\frac{\phi(0)}{\phi(x)}} \right) f^2(x) dx \right)^2 - F_2(s, t) \right\}^{1/2} \\
&\quad \times \int_0^1 u g^2(u) du - F_1(s, t). \tag{14}
\end{aligned}$$

Since $g(u) \neq 0$ for all $u \in (0, 1)$, we have $f(x) \neq 0$. Hence $F_1(s, t) > 0$ and $F_2(s, t) > 0$. Thus, theorem is proved. \square

Remark 3.1. For $\phi(x) = \psi(x) = x$, inequality (13) reduces to inequality (13) proved by Gao [2]. Hence inequality (13) is a generalization of the corresponding result of Gao [2].

Remark 3.2. If $F_1(s, t)$ and $F_2(s, t)$ contained in (13) are replaced by zero simultaneously, then inequality (13) reduces to the following form:

$$\int_0^\infty f^2(x) dx < \frac{1}{(\inf\{\phi'(x)\})^2} \int_0^1 u g^2(u) du.$$

Clearly, this is also a generalization of the Hardy–Littlewood inequality (see Hardy et al. [1]).

References

- [1] G.H. Hardy, J.E. Littlewood, G. Polya, *Inequalities*, Cambridge Univ. Press, London, 1952.
- [2] Mingzhe Gao, On Hilbert's integral inequality, *Math. Appl.* 11 (3) (1998) 32–35.